



Managing children who are sick, infectious, or with allergies

Policy statement

We aim to provide care for healthy children through preventing cross infection of viruses and bacterial infections and promote health through identifying allergies and preventing contact with the allergenic trigger.

Procedures for children who are sick or infectious

- If a child appears unwell during the day, for example has a raised temperature, sickness, diarrhoea* and/or pains, particularly in the head or stomach then the setting manager calls the parents and asks them to collect the child or send a known carer to collect on their behalf. *Diarrhoea is defined as 3 or more liquid or semi-liquid stools in a 24-hour period. www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities/chapter-9-managing-specific-infectious-diseases#diarrhoea-and-vomiting-gastroenteritis
- If a child has a raised temperature, they are kept cool by removing top clothing, giving cold water to drink and kept away from draughts.
- A child's temperature is taken and checked regularly, using digital thermometer kept on hook by first aid kits.
- If the child's temperature does not go down and is high, then we may give them Calpol, after first obtaining verbal consent from the parent where possible and on the understanding, they are on their way to collect their child. This will only take place in exceptional circumstances as the use of emergency medicine does not apply to children over two years of age and we are not allowed to 'prescribe'. Parents sign the medication record when they collect their child.
- In an emergency an ambulance is called and the parents are informed.
- Parents are advised to seek medical advice before returning them to the setting; the setting can refuse admittance to children who have a raised temperature, sickness and diarrhoea or a contagious infection or disease.
- Where children have been prescribed antibiotics for an infectious illness or complaint, parents are asked to keep them at home for 48 hours.
- After diarrhoea or vomiting, parents are asked to keep children home for 48 hours following the last episode.
- Where children have had a raised temperature, they should stay at home until they have been fever-free for 24 hours, without the use of fever-reducing medication.
- Some activities such as sand and water play and self-serve snack will be suspended for the duration of any outbreak.
- The setting has information about excludable diseases and exclusion times. The full list is obtainable from https://www.publichealth.hscni.net/sites/default/files/Guidance_on_infection_control_in%20schools_poster.pdf
- The setting manager notifies the committee if there is an outbreak of an infection (affects more than 3-4 children) and keeps a record of the numbers and duration of each event.
- The setting manager has a list of notifiable diseases, which does include COVID-19 (see Health Folder) and contacts Public Health England (PHE) and Ofsted in the event of an outbreak.
- If staff suspect that a child who falls ill whilst in their care is suffering from a serious disease that may have been contracted abroad such as Ebola, immediate medical assessment is required. The setting manager or deputy calls NHS111 and informs parents.

- It should also be noted the setting cannot take bottles of non-prescription medicine from parents to hold on a 'just in case' basis, unless there is an immediate reason for doing so.

HIV/AIDS procedure

HIV virus, like other viruses such as Hepatitis, (A, B and C), are spread through body fluids. Hygiene precautions for dealing with body fluids are the same for all children and adults.

- Single use vinyl gloves and aprons are worn when changing children's nappies, pants and clothing that are soiled with blood, urine, faeces or vomit.
- Protective rubber gloves are used for cleaning/sluicing clothing after changing.
- Soiled clothing is rinsed and bagged for parents to collect.
- Spills of blood, urine, faeces or vomit are cleared using mild disinfectant solution and mops; cloths used are disposed of with clinical waste.
- Tables and other furniture or toys affected by blood, urine, faeces or vomit are cleaned using a disinfectant.
- Baby mouthing toys are kept clean and plastic toys cleaned in sterilising solution regularly.

Nits and head lice

- Nits and head lice are not an excludable condition; although in exceptional cases parents may be asked to keep the child away from the setting until the infestation has cleared.
- On identifying cases of head lice, all parents are informed and asked to treat their child and all the family, using current recommended treatments methods if they are found.

Further guidance

For more information on infectious diseases see following link www.gov.uk/government/publications/health-protection-in-schools-and-other-childcare-facilities

Procedures for children with allergies

- When children start at the setting we ask their parents if their child suffers from any known allergies. This is recorded on the Registration Form.
- If a child has an allergy, we complete an allergy risk assessment form to detail the following:
 - the risk identified – the allergen (i.e. the substance, material or living creature the child is allergic to such as nuts, eggs, bee stings, cats etc.)
 - the level of risk, taking into consideration the likelihood of the child coming into contact with the allergen
 - control measures, such as prevention from contact with the allergen
 - review measures
- A Health Care plan will also be completed which will include
 - the nature of the reaction e.g. anaphylactic shock reaction, including rash, reddening of skin, swelling, breathing problems etc.
 - managing allergic reactions, medication used and method (e.g. EpiPen)
- This risk assessment form and Health Care Plan is kept in the child's personal file or with the child's medication and is shared with all staff.
- The child's name is added to the Allergies/Dietary Requirements list.
- Parents show staff how to administer medication in the event of an allergic reaction.
- No nuts or nut products are used within the setting.
- Parents are made aware, so that no nut or nut products are accidentally brought in.
- Any foods containing food allergens are identified on children's menus.

Oral Medication

- Oral medication must be prescribed or have manufacturer's instructions written on them.
- Staff must be provided with clear written instructions for administering such medication.
- All risk assessment procedures are adhered to for the correct storage and administration of the medication.
- The setting must have the parents' prior written consent. Consent is kept on file.

Life-saving medication and invasive treatments

Life-saving medication and invasive treatments may include adrenaline injections (Epi pens) for anaphylactic shock reactions (caused by allergies to nuts, eggs etc) or invasive treatment such as rectal administration of Diazepam (for epilepsy).

- The key person responsible for the intimate care of children who require life-saving medication or invasive treatment will undertake their duties in a professional manner having due regard to the procedures listed above.
- The child's welfare is paramount, and their experience of intimate and personal care should be positive. Every child is treated as an individual and care is given gently and sensitively; no child should be attended to in a way that causes distress or pain.
- The key person works in close partnership with parents/carers and other professionals to share information and provide continuity of care.
- Children with complex and/or long-term health conditions have a health care plan (04.2a) in place which takes into account the principles and best practice guidance given here.
- Key persons have appropriate training for administration of treatment and are aware of infection control best practice, for example, using personal protective equipment (PPE).
- Key persons speak directly to the child, explaining what they are doing as appropriate to the child's age and level of comprehension.
- Children's right to privacy and modesty is respected. Another practitioner is usually present during the process.

Record keeping

For a child who requires invasive treatment the following must be in place from the outset:

- a letter from the child's GP/consultant stating the child's condition and what medication if any is to be administered
- written consent from parents allowing members of staff to administer medication
- proof of training in the administration of such medication by the child's GP, a district nurse, children's nurse specialist or a community paediatric nurse
- an Individual Healthcare Plan

Copies of all letters relating to these children must be sent to the insurance provider for appraisal. Confirmation will then be issued in writing confirming that the insurance has been extended. If we are unsure about any aspect, we will contact the Morton Michel Tel 0330 058 9861 (9.00am – 5.00pm Mon – Fri) or e-mail customer.service@mortonmichel.com

A record is made in the medication record book of the intimate/invasive treatment each time it is given.

Physiotherapy

- Children who require physiotherapy whilst attending the setting should have this carried out by a trained physiotherapist.
- If it is agreed in the health care plan that the key person should undertake part of the physiotherapy regime then the required technique must be demonstrated by the physiotherapist personally; written guidance must also be given and reviewed regularly. The physiotherapist should observe the practitioner applying the technique in the first instance.

Safeguarding/child protection

- Practitioners recognise that children with SEND are particularly vulnerable to all types of abuse, therefore the safeguarding procedures are followed rigorously.
- If a practitioner has any concerns about physical changes noted during a procedure, for example unexplained marks or bruising then the concerns are discussed with the designated person for safeguarding and the relevant procedure is followed.

Treatments such as inhalers or Epi-pens must be immediately accessible in an emergency.

This policy should also be read in conjunction with our Administering Medicines policy.